

UNIT III – APPLICATIONS OF PARTIAL DIFFERENTIAL EQUATIONS**PART -A**

1. Classify the partial differential equation $4 \frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$

ANS

Here A=4 ; B=0; C=0

$$B^2-4AC = 0-0=0$$

The PDE is parabolic

2. In the equation of motion of vibrating string $a^2 \frac{\partial^2 y}{\partial x^2} = \frac{\partial^2 y}{\partial t^2}$, what does a^2 stand for?

ANS

$a^2 = \frac{T}{m}$ where T is the tension of the string and m is the mass per unit length of the string

3. Write down all possible solutions of one dimensional wave equation

ANS

$$\begin{aligned} \oplus y(x,t) &= [Ae^{px} + Be^{-px}][Ce^{pat} + De^{-pat}] \\ \oplus y(x,t) &= [A\cos(px) + B\sin(px)][C\cos(pat) + D\sin(pat)] \\ \oplus y(x,t) &= [Ax + B][Ct + D] \end{aligned}$$

4. A tightly stretched string with fixed end points $x=0$ and $x=l$ is initially in a position given by

$y(x,0) = y_0 \sin^3\left(\frac{\pi x}{l}\right)$. If it is released from rest in this position, write the boundary conditions.

ANS

The boundary conditions are

$$\times y(0,t) = 0$$

$$\times y(l,t) = 0$$

$$\times \left(\frac{\partial y}{\partial t}\right)_{t=0} = 0$$

$$\times y(x,0) = y_0 \sin^3\left(\frac{\pi x}{l}\right)$$

5. If ends of a string of length 'l' are fixed and the midpoint of the string is drawn aside through a height 'h' and the string is released from rest, state the initial and boundary conditions.

ANS

The conditions are

$$\bullet y(0,t) = 0$$

$$\bullet y(l,t) = 0$$

$$\bullet \left(\frac{\partial y}{\partial t}\right)_{t=0} = 0$$

$$y(x,0) = \begin{cases} \frac{2hx}{l}, & 0 \leq x \leq l/2 \\ \frac{2h(l-x)}{l}, & l/2 \leq x \leq l \end{cases}$$

6. Write the initial conditions of the wave equation if the string has an initial displacement $f(x)$ but no initial velocity

ANS

The displacement $y(x,t)$ is from $a^2 \frac{\partial^2 y}{\partial x^2} = \frac{\partial^2 y}{\partial t^2}$

The conditions are

$$\oplus y(0,t) = 0$$

$$\oplus y(l,t) = 0$$

$$\oplus \left(\frac{\partial y}{\partial t} \right)_{t=0} = 0$$

$$\oplus y(x,0) = f(x)$$

7. Write the initial conditions of the wave equation if the string has an initial velocity $g(x)$ but has no initial displacement.

ANS

The displacement $y(x,t)$ is from $a^2 \frac{\partial^2 y}{\partial x^2} = \frac{\partial^2 y}{\partial t^2}$

The conditions are

$$\oplus y(0,t) = 0$$

$$\oplus y(l,t) = 0$$

$$\oplus y(x,0) = 0$$

$$\oplus \left(\frac{\partial y}{\partial t} \right)_{t=0} = g(x)$$

8. What does a^2 represent in one dimensional heat flow equation $u_t = a^2 u_{xx}$?

ANS

$a^2 = \frac{k}{pc}$ where k is the thermal conductivity; c is the density and p is the specific heat

9. In steady state conditions, derive the solution of one dimensional heat flow equation

ANS

The one dimensional heat flow equation is $\alpha^2 \frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$

In steady state, $\frac{\partial u}{\partial t} = 0$

$$\Rightarrow \frac{\partial^2 u}{\partial x^2} = 0$$

$$\Rightarrow \frac{\partial u}{\partial x} = a$$

$$\Rightarrow u = ax + b$$

10. Write down the three possible solutions of one dimensional heat equation

ANS

- ✓ $u(x,t) = [Ae^{px} + Be^{-px}] Ce^{\alpha^2 p^2 t}$
- ✓ $u(x,t) = [A \cos(px) + B \sin(px)] Ce^{-\alpha^2 p^2 t}$
- ✓ $u(x,t) = [Ax + B]C$.

11. What is the basic difference between the solution of one dimensional wave equation and one dimensional heat equation?

ANS

| S.NO. | One dimensional wave equation | One dimensional heat equation |
|-------|--|--|
| 1 | $a^2 \frac{\partial^2 y}{\partial x^2} = \frac{\partial^2 y}{\partial t^2}$ is hyperbolic | $\alpha^2 \frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$ is parabolic |
| 2 | The suitable solution $y(x,t) = [A \cos px + B \sin px][C \cos pt + D \sin pt]$ is periodic w.r.t.time | The suitable solution $u(x,t) = [A \cos(px) + B \sin(px)]C e^{-\alpha^2 p^2 t}$ is not periodic with respect to time |

12. What are the possible solutions for Laplace equation $U_{xx} + U_{yy} = 0$ by method of separation of variables? (OR) Write all the three possible solutions of steady state two dimensional heat equation

ANS

- $u(x,y) = [Ae^{px} + Be^{-px}][C \cos(py) + D \sin(py)]$
- $u(x,y) = [A \cos(px) + B \sin(px)][Ce^{py} + De^{-py}]$
- $u(x,y) = [Ax + B][Cy + D]$

PART – B

1. A uniform string is stretched and fastened to two points ‘l’ apart. Motion is started by displacing the string into the form of the curve $y=kx(l-x)$ and then released from this position at time $t=0$. Derive the expression for the displacement of any point of the string at a distance x from one end at time t .
2. A tightly stretched string with fixed end points $x=0$ and $x=l$ is initially displaced in the position $y = y_0 \sin^3\left(\frac{\pi x}{l}\right)$ and then released from rest. Find the displacement y at any distance x from one end at time t .
3. A tightly stretched string of length ‘l’ has its ends fastened at $x=0$ and $x=l$. The midpoint of the string is then taken to height h and released from rest in that position. Find the displacement of a point of the string at time t from the instant of release.
4. A tightly stretched string of length l is initially at rest in its equilibrium position and each of its points is given the velocity $V_0 \sin^3\left(\frac{\pi x}{l}\right)$. Find the displacement $y(x,t)$.
5. A string of length l is initially at rest in its equilibrium position and motion is started by giving each of the points a velocity given by $v = \begin{cases} cx, 0 \leq x \leq l/2 \\ c(l-x), l/2 \leq x \leq l \end{cases}$. Find the displacement function $y(x,t)$.

6. A rod 30 cm long has its ends A and B kept at 30°C and 80°C respectively, until steady state conditions prevail. The temperature at each end is then suddenly reduced to 0°C and kept so on. Find the resulting temperature function $u(x,t)$ taking $x=0$ at A.
7. The ends A and B of a rod 40 cm long have their temperature kept at 0°C and 80°C respectively until steady state condition prevails. The temperature of the end B is then suddenly reduced to 40°C and kept so while that of the end A is kept at 0°C . Find the subsequent temperature distribution $u(x,t)$ in the rod.
8. A square plate is bounded by the lines $x=0$; $y=0$; $x=20$ and $y=20$. Its faces are insulated. The temperature along the upper horizontal edge is given by $u(x,20) = x(20-x), 0 < x < 20$, while the other edges are kept at 0°C . Find the steady state temperature distribution in the plate.
9. A rectangular plate with insulated surfaces is 20 cm wide and so long compared to its width that it may be considered infinite in length. If the temperature at the short edge $x=0$ is given by $u = \begin{cases} 10y, 0 \leq y \leq 10 \\ 10(20-y), 10 \leq y \leq 20 \end{cases}$ and the two long edges as well as the other short edge are kept at 0°C . Find the steady state temperature distribution in the plate.
10. A rectangular plate with insulated surfaces is 10 cm wide and so long compared to its width that it may be considering infinite in length. The temperature at short edge $y=0$ is given by $u = \begin{cases} 20x, 0 \leq x \leq 5 \\ 20(10-x), 5 \leq x \leq 10 \end{cases}$ and all the other edges are kept at 0°C . Find the steady state temperature.

UNIT-III-APPLICATION OF PARTIAL DIFFERENTIAL EQUATIONS

PART-A

1. Classify the P.D.E $\frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$.
2. Write down all the possible solution of one dimensional wave equation.
3. Write all three possible solutions of steady state two dimensional heat equations.
4. A rod 40cm long insulated sides has its ends A and B kept at 20°C and 60°C respectively. Find the steady state temperature at a location 15cm from A
5. A tightly stretched string with fixed end points $x=0$ and $x=l$ is initially in a position given by $y(x,0) = y_0 \sin^3\left(\frac{\pi x}{l}\right)$, if it is released from rest in its position, write the boundary conditions.
6. What does a^2 represents in one dimensional heat flow equation $u_t = a^2 u_{xx}$.
7. A rod 20cm long insulated sides has its ends A and B kept at 30°C and 90°C respectively. Find the steady state temperature distribution on the rod.
8. Write down all the possible solution of one dimensional heat equation.
9. The end of a string of length l is fixed at both ends. The midpoint of the string is taken to a height λ and then released from rest. Write the initial conditions of the string.
10. Classify the P.D.E $y^2 U_{xx} - 2xy U_{xy} + x^2 U_{yy} + 2U_x - 3U_y = 0$.

PART-B

1. A string is stretched and fastened to two points l apart motion is stretched by displacing the string into the form $y=k(l/x-x^2)$ and then released it from the position at time $t=0$. Find the displacement of the point of the string at a distance of x from one end at time t .
2. A tightly stretched string of length $2l$ has its end fastened at $x=0$ & $x=2l$. The midpoint of the string is then taken to height b and released from rest in that position. Find the lateral displacement of a point of the string at time t from the instant of release.
3. A string of length l is initially at rest in equilibrium position and each of its point is given the velocity $V_0 \sin\left(\frac{\pi x}{l}\right)$ $0 < x < l$. Compute the displacement of the string.

4. A square plate is bounded by the lines $x=0, y=0, x=20$ and $y=20$. Its faces are insulated. The temperature along the upper horizontal edge is given by $u(x, 20) = x(20 - x), 0 < x < 20$ while the other two edges are kept at 0°C . Find the steady state temperature distribution in the plate.
5. A tightly stretched string of length l has its end fastened at $x=0$ and $x=l$. The midpoint of the string is then taken to height b and released from rest in that position. Find the lateral displacement of a point of the string at time t from the instant of release.
6. A rectangular plate with insulated surface is 10 cm wide and so long compared to its width that it may be considered infinite in length without introducing appreciable error. The temperature at short edge $y=0$ is given by $u = \begin{cases} 20x, 0 \leq x \leq 5 \\ 20(10 - x), 5 \leq x \leq 10 \end{cases}$ and all the other three edges are kept at 0°C . Find the steady state temperature at any point in the plate.
7. A tightly stretched string with fixed end points $x=0$ and $x=l$ is initially at rest in its equilibrium position. If it is set vibrating giving each point a initial velocity $3x(l-x)$, find the displacement.
8. A rod 30cm long has its end A and B kept at 20°C and 80°C respectively, until steady state conditions prevail. The temperature at each end is then suddenly reduced to 0°C and kept so. Find the resulting temperature distribution function $u(x, t)$ taking $x=0$ at A.
9. If a string l is initially at rest in its equilibrium position and each of its points is given a velocity v such that $v = \begin{cases} kx, 0 < x \leq \frac{\ell}{2} \\ k(\ell - x), \frac{\ell}{2} < x \leq \ell \end{cases}$. Find the displacement $y(x, t)$.
10. A rod of length l has its end A and B kept at 0°C and 100°C respectively until steady state condition prevails. If the temperature at B is reduced suddenly to 0°C and kept so while that of A is maintained find the temperature $u(x, t)$ at a distance x from A and at time t .